



## Abstract

Eutrophication is a major issue for many bodies of water and ~~coasts~~ around the world.

Phytoextraction has been used as a tool to mitigate eutrophication in the past, however it has not been done with agricultural crops to produce a harvestable product while removing nutrients from stream systems. In this study we ~~used~~ eddy systems and used watercress plants to mitigate nutrient loading in a closed system. We found that the watercress removed phosphate from the system but was ineffective at removing ammonium and nitrate in our modelled system. Future studies are ~~needed~~ to find a reduction in nitrogen levels and implement this method in natural streams that experience nutrient loading.

Keywords:

Highly impacted landscapes, such as those where small streams and tributaries to larger rivers, are influenced by proximity of intense agricultural or urbanization. Once excess nutrients reach lakes and estuaries, the ecological effects magnify and the effects are much harder to mitigate. Literature does not address the intermediate phase between the source pollution and the larger stagnant body of water. In addition, few studies have examined the high levels of toxin buildup in flowing systems such as streams, and only focus on stagnant waters. Therefore, it is necessary to consider the impacts of concentrated nutrient runoff into stream systems before they contaminate larger ecosystems.

One potential means of reducing surface water nutrient concentrations is through phytoremediation. Phytoremediation is the use of plants to absorb pollutants from an ecosystem and can be used to treat eutrophic systems with plants that absorb high concentrations of N and P (Király et al., 2013; Lu, 2010). Elevated N levels produce algae blooms that compete with fish and other aquatic species at high biomass (Chen et al., 2009); in removing excess N from the system, these ecosystems will be more productive. Similar ideas have been implemented to mitigate contaminants in pond and lake systems using inedible tropical plants (Todd, 2016). Phytoremediation has also been implemented on land, where plants are used to absorb contaminants such as lead or other heavy metals (Salt et al., 1995).

Although phytoremediation in point source meshes and storm water ponds has been previously examined (Hunt, 2006), flowing surface waters such as small streams and rivers have received less attention. Eddy flow systems are of particular interest because of longer retention

Torrance, 1973; Ensign & Doyle, 2006). Thus, eddies provide conditions for plants to extract excess N and P that cause eutrophication (Zhao et al., 2012; Schnoor et al., 1995).

We proposed an experiment to evaluate stream eddy nutrient uptake by harvestable watercress. We hypothesized that nutrient concentrations will decrease as plant uptake increases, until a nutrient becomes limiting (Mengel & Kirkby, 1978; Tanner, 1996). We also hypothesized that a slow moving eddy system will foster nutrient uptake (Carpenter et al., 1998). We studied uptake rate in modeled eddy systems to examine the effects of plant traits and tissue stoichiometry on phytoremediation efficiency. For example, watercress grows rapidly and is known to uptake more nutrients per unit tissue than species in the *brassicaceae* family (Tanner, 1996). Watercress is also a very marketable agricultural crop that has the potential to increase the feasibility of implementing it in eddy systems. These experiments allow us to evaluate an effective method to reduce excess nutrient loading in surface waters that is cost and resource efficient. Further research could investigate the effective uptake of harmful organic and inorganic waste for selective uptake (McKone & Maddalena, 2007).

## Methods

### Eddy modeling

To uptake excess nutrients in a stream system, we modeled eddies in Skidmore's greenhouse. Eddies generally have long retention times and lower velocities in streams, which allows plants more time to extract excess nitrogen and phosphorus from the water that causes eutrophication further downstream into lentic ecosystems (Pauer & Auer, 2000). In these lake and pond ecosystems, stratification makes it more difficult to use phytoextraction as a method of mitigating eutrophication.

We modeled eddys using three rectangular basins with a 5.258 x 10<sup>-3</sup> m<sup>3</sup>/s pump in each one to keep the water moving with an average velocity of 0.03 m/s circular motion. We also added an airstone from an aerator to each basin to eliminate anoxic conditions and promote aerobic bacteria, important for nutrient uptake in stream ecosystems. The outer bottom half of each basin was wrapped in black plastic to prevent much light from harming the plant roots from the side of the container, as solar radiation would not be natural at this angle in stream.

The plants were placed in a raft made of condensed Styrofoam insulation board with a reflective surface on the outside. Holes were cut into the insulation board to accommodate 5.08 cm net pots, spaced about 7.62 cm apart. Using this method, the plants can expand their roots into the water and absorb nutrients easily. Each of the three Styrofoam rafts holds 24 net pots. We placed the plants in the rafts to grow their root systems and above ground biomass, four weeks before we began the study.

## Plant Growth

We started the watercress plants with stem cuttings from a nearby stream. We placed a five cm cutting into each Rockwool cube of a 98 cube perforated tray. We placed these Rockwool trays into a shallow tray containing our compost tea nutrient solution diluted 50 %. To grow out our plants quickly, we installed a 60-watt T8 fluorescent grow light fixture 20 cm above the tops of the Rockwool trays. These lights were set for a 24 hr period to promote rapid growth. Once the watercress was well rooted (12 days), we transferred each plant to a 5 cm net pots and placed them into a floating raft system with the same nutrient solution and under the same

lighting conditions to have the roots extend and cover more surface area in the water column. This allowed the watercress to uptake as much nitrogen and phosphorus as possible. Once the watercress cuttings were well rooted and growing in the floating raft system for two weeks, we

## Nutrient Concentrations and Sources

For nutrients, we made a solution of compost tea that consisted of 4.500 kg of composted cow manure and plant material mixed with 30 liters of water. We left this mixture for 48 hours with a pump to keep it aerated and mixing before we filtered it all through a strainer and coffee filter to get our concentrated nutrient solution. We diluted this compost tea solution to 50 % in tap water to grow the watercress before the study began, and as the organic fertilizer experimental addition.

For our synthetic fertilizer trial, we added Age Old Grow hydroponic fertilizer at a rate of 1/2 tablespoon per gallon of water. Age Old Grow contains 12 % total nitrogen (3 % water insoluble





The data show a decreasing trend in phosphate using both organic and inorganic fertilizers, as well as a high increase in nitrate with the organic fertilizer and high increase in ammonium with the inorganic fertilizer.

released due to mineralization within the basin as the particles may not have been completely dissolved especially near the root mat of the watercress (Westerman & Tucker, 1974). Another reason may be that the growth of cyanobacterial along the edges of the basins and top of the rockwool cubes could have fixed more nitrogen from the atmosphere and into the water (Vitousek et al., 2002). Finally, there was slight evaporation from the bins, as well as potential stress and die-off from plant roots that could have made the later samples more concentrated.

Future studies could also include changing the ratio of plants to the surface area or volume of the eddy to see what effect it may have on the nutrient uptake rate and amount of harvestable watercress that can be grown. Higher densities of watercress may produce higher harvestable yield while taking up more nutrients.

To examine the model more specifically, we could measure nutrient uptake in the plant tissue instead of the surrounding water (Mattina et al., 2003; Cataldo et al., 1975). We could also trace the isotopes throughout the modelled eddy system to determine the fate of the nutrients and how much is being stored in bacteria as well as the pace of the enzyme activity (David & Raven, 1992). A closer analysis of these processes helps us understand how the model can best represent a natural system, and where to channel effective nutrient absorption to prevent eutrophication and produce a harvestable crop.

Through experimentation with using eddys to grow harvestable watercress while mitigating nutrient loading in stream systems, we have found a significant decrease in phosphate concentrations. This means that farming eddys would be feasible as a method of mitigating nutrient loading in phosphorus limited lake systems. Our results did not show a decrease in nitrogen concentrations in the form of ammonium and nitrate, the (r)3 (a)4 (t)-2 (e)4 (, t)>(v)2 tts(t)-2 (e

## References



Schaller, J. L., Royer, T. V., David, M. B., & Tank, J. L. (2004). Denitrification associated with plants and sediments in an agricultural stream. *Journal of the North American Benthological Society*, 23(4), 667-676.

Schnoor, J. L., Light, L. A., McCutcheon, S. C., Wolfe, N. L., & Carreira, L. H. (1995). Phytoremediation of organic and inorganic contaminants. *Environmental Science & Technology*, 29(7), 318A-323A.

Smith, V. H., Tilman, G. D., & Nekola, J. C. (1999). Eutrophication: impacts of excess nutrient